


## Can cells see each other?

### **What could be the biological function of cellular infrared 'vision'?**

As shown in the flow diagram below, our work up to this point seems to link cellular infrared 'vision' with animal development which depends critically on cell migration .




Therefore, we suspect that the infrared 'vision' of cells plays a role in development.

There is an additional reason for this conjecture. Animal cell migration is very slow and animal development occurs simultaneously everywhere in the embryo. What, if a migrating cell makes a mistake and ends up in the wrong place? It cannot stop the development everywhere else, turn back the developmental clock and start from scratch. Therefore, it seems good engineering to provide the migrating cells with the means to 'look ahead' towards their spatial goals.

How far should a cell look ahead? In a developing embryo any spatial goal is a moving target. If it takes a cell 6-10 hours to reach this goal, it would probably no longer exist because development may have changed the area around the goal too much. Since an animal cell needs about 1.5 hours to cross its own diameter of appr. 20  $\mu\text{m}$  its spatial goals should be no further away than 4-6 cell diameters or 80-120  $\mu\text{m}$ . Indeed, our experiments showed that the cells ignored microscopic light sources that were further away than about 60  $\mu\text{m}$ .

### **Cells detect each other at a distance.**


It stands to reason that the major kinds of spatial goals of cells in an embryo are other cells. For many years I observed that approaching cells reached out towards each other from a distance, as if they 'saw' each other. The figure below shows a remarkable example.




In this case an epithelial cell from an African Green Monkey (upper left corner) extends in a very targeted way towards a fibroblast from a mouse that had turned its front towards the epithelial cell at an earlier time. In spite of the differences of animal and tissue type, the 2 cells recognized their presence at a distance and actively approached each other.

**Cells react to each other across a film of glass.**

In order to test whether such cell-to-cell encounters were mediated by some chemicals, I separated cells by a thin, but chemically impenetrable glass film. The figure below shows a Hamster fibroblast sitting on one side of the glass film (Let us call it the 'A'-face).



On the other side of the glass film (Let us call it the 'B'-face) we forced cells to grow in a pattern of parallel stripes (see green lines on the figure below).



We found that the cells on the 'A'-face oriented themselves predominantly perpendicular to the cell stripes on the 'B'-face (see red lines on the above figure); even though the 2 faces were insulated from each other by the impenetrable and inert glass film. When we coated the 'A'-face with an opaque gold film and then tried the experiment, no such special orientation was found. On the other hand, if we coated it with a film of silicone which is opaque for visible light and transparent for near infrared light, the special orientation of the cells reappeared. In other words, only as long as infrared light could pass through the glass film, the cells were also able to determine the direction of the cell stripes on the opposite face.

### **Significance for cell intelligence:**

#### *Infrared 'vision' may represent a novel form of cell-to-cell communication*

The observation of a perpendicular orientation between cells in different planes is not the amazing results. Many tissues in the body (e.g. the cornea, the intestines and many others) are weaved like this. Cells seem to be programmed to line up side-by-side in the same plane and perpendicular to each other in the planes above and below. The amazing result is the observation that the cells on the 'A'-face knew the orientation of the cells on the 'B'-face if near-infrared light could pass across the glass film. Similar to the argument about cells following [guiding 'roads'](#) the cells on the 'A'-face had to 'see' at least 2 different points on the cell stripes on the other side of the glass film in order to determine the direction of the stripes. In the above experiment, however, we did not provide any pulsating infrared light. Therefore, if the cells themselves were the sources of the light that passed through the glass, they were communicating their spatial orientation to the cells in a layer above. Therefore, it seems possible that the infrared 'vision' of cells is used in the developing animal to weave tissues.

### **Source:**

<http://www.basic.northwestern.edu/g-buehler/film.htm>

## **Bibliography**

### **(Selected articles)**

1. Albrecht-Buehler, G.: Daughter 3T3 cells: Are they mirror images of each other? J. Cell Biol. 72: 595-603, 1977
2. Albrecht-Buehler, G.: The phagokinetic tracks of 3T3 cells. Cell 11:395-404, 1977
3. Albrecht-Buehler, G.: The phagokinetic tracks of 3T3 cells: Parallels between the orientation of track segments and of cellular structures which contain actin or tubulin. Cell 12:333-339, 1977

4. Albrecht-Buehler, G.: The tracks of moving cells. *Scientific American* 238:68-76, 1978
5. Albrecht-Buehler, G.: The angular distribution of directional changes of guided 3T3 cells. *J. Cell Biol.* 80:53-60, 1979
6. Albrecht-Buehler, G. and Bushnell, A.: The orientation of centrioles in migrating 3T3 cells. *Exp. Cell Res.* 120: 111-118, 1979
7. Albrecht-Buehler, G.: Group locomotion of PtK1 cells. *Exp. Cell Res.* 122:402-407, 1979
8. Albrecht-Buehler, G.: The autonomous movements of cytoplasmic fragments. *Proc. Natl. Acad. Sci. U.S.A.* 77: 6639-6644, 1980
9. Albrecht-Buehler, G.: Does the geometric design of centrioles imply their function? *Cell Motility* 1: 237-265, 1981
10. Albrecht-Buehler, G. Is Cytoplasm Intelligent too? In: *Muscle and Cell Motility VI* (ed. J. Shay) p. 1-21 (1985).
11. Albrecht-Buehler, G. In defense of non-molecular' cell biology. *International Review of Cytology* 120:191-241 (1990)
12. Albrecht-Buehler, G. The iris diaphragm Model of centriole and basal body formation. *Cell Motility and the Cytoskeleton* 17:187-213 (1990)
13. Albrecht-Buehler, G. Surface extensions of 3T3 cells towards distant infrared sources. *J. Cell Biol.* (1991)114:493-502
14. Albrecht-Buehler, G. (1992) Speculation about the function and formation of centrioles and basal bodies. In: *The Centrosome* (ed. V.I. Kalnins) Academic Press, pp 69-102
15. Albrecht-Buehler, G. A rudimentary form of cellular 'vision'(1992) *Proc. Natl. Acad. Sci. USA* 89:8288-8292
16. Albrecht-Buehler, G. The cellular infrared detector appears to be contained in the centrosome. *Cell Motility and the Cytoskeleton* 27:262-271 (1994)
17. Albrecht-Buehler, G. Changes of cell behavior by near-infrared signals. *Cell Motility and the Cytoskeleton* 32:299-304 (1995).
18. Albrecht-Buehler, G., Autofluorescence of live purple bacteria in the near infrared. (1997) *Experimental Cell Research* 236:43-50
19. Albrecht-Buehler, G. Phagokinetic Track Assay of Cell locomotion in Tissue Culture, In 'Cells: A Laboratory Manual' Vol. 2, Cold Spring Harbor, NY (1997) 77.1-77.10

20. Albrecht-Buehler, G. (1998) Altered drug resistance of microtubules in cells exposed to infrared light pulses: Are Microtubules the 'Nerves' of Cells? *Cell Motility and the Cytoskeleton* 40:183-192

21. Albrecht-Buehler, G. (2005) A Long-Range Attraction Between Aggregating 3T3 Cells Mediated By Near-Infrared Light Scattering, *PNAS* 102 no 14: 5050-5055 (<http://www.pnas.org/cgi/reprint/0407763102v1>)